

ARMY RESEARCH LABORATORY



Nonmonoenergetic Beam With Helical Čerenkov Radiation: Possible Technique to Determine Magnetic Field in a Hostile Electromagnetic Environment

Josip Z. Šoln

ARL-TR-947

January 1996

19960213 069

DTIC QUALITY INSPECTED 4

NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 1996		3. REPORT TYPE AND DATES COVERED Final, Oct 1994-Jun 1995
4. TITLE AND SUBTITLE Nonmonoenergetic Beam With Helical Čerenkov Radiation: Possible Technique to Determine Magnetic Field in a Hostile Electromagnetic Environment			5. FUNDING NUMBERS PN: 1L162618AH80	
6. AUTHOR(S) Josip Z. Šoln				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: AMSRL-SL-CM Aberdeen Proving Ground, MD 21010-5423			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-947	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Modifications caused by the nonmonoenergetic beams to the recently described helical Čerenkov radiation (resulting from electron helical motion in a medium under the influence of magnetic field) are studied in a visible portion of the spectrum. The analysis is greatly simplified by utilizing a new analytical approximate expression for the number of emitted photons per unit path length for the usual "monoenergetic" helical Čerenkov radiator. The effect of the nonmonoenergetic beam is that simultaneously with the helical Čerenkov radiation also the radiation into harmonics (above and below the helical Čerenkov effect threshold) may occur at the same radiation angle and at the same visible frequency. Hence, harmonic radiators may enhance the usual helical Čerenkov radiation. For the medium of silica aerogel with the index of refraction of 1.075 and the beam energy in the 2-3-MeV range interacting with the magnetic field of about 10 T, one estimates that in the visible portion of the spectrum an electron from a nonmonoenergetic beam will generate between 1 and 2 photons at the end of the 10-cm interaction length. This shows the possibility of detecting a magnetic field coming from an electromagnetic treat wave form, which in turn, represents a hostile electromagnetic environment. The importance of detecting this magnetic field comes from the fact that in an environmental medium it can disrupt optical signals. Here we show the way for measuring and quantifying such disruptions.</p>				
14. SUBJECT TERMS helical Čerenkov effect, dielectric medium, harmonics			15. NUMBER OF PAGES 16	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

INTENTIONALLY LEFT BLANK.

TABLE OF CONTENTS

	<u>Page</u>
1. POWER SPECTRA FOR THE HELICAL ČERENKOV EFFECT AND HARMONIC RADIATION ABOVE AND BELOW HELICAL ČERENKOV THRESHOLD	1
2. NUMBER OF PHOTONS PER UNIT PATH LENGTH IN THE VISIBLE SPECTRUM	2
3. APPLICATIONS	7
4. REFERENCES	11
DISTRIBUTION LIST	13

INTENTIONALLY LEFT BLANK.

1. POWER SPECTRA FOR THE HELICAL ČERENKOV EFFECT AND HARMONIC RADIATION ABOVE AND BELOW HELICAL ČERENKOV THRESHOLD

The helical Čerenkov effect (HCE) arises from electrons moving in a medium on helical trajectories under the influence of the uniform magnetic field that, for simplicity, is defined as $\vec{B} = \hat{z}B$, with B denoting the magnitude of the magnetic field and \hat{z} its direction (Šoln 1992). The helical Čerenkov radiation with angular frequency ω will occur if (Šoln 1992)

$$n(\omega)\beta_z(0) \geq 1, \quad (1)$$

where $\beta_z(0) = v_z(0)/c$, with $v_z(0)$ denoting the parallel component of the electron velocity with respect to \vec{B} . Here and in what follows, $\beta = v/c$, $\beta_z = v_z/c$, $\beta_\perp = v_\perp/c$ with v_\perp being the perpendicular component of the electron velocity with respect to \vec{B} and $v^2 = v_z^2 + v_\perp^2$. One notices that v_z is also the velocity of the electron-guiding center. The power spectra for the HCE and for the harmonic radiation below and above the helical Čerenkov threshold are, respectively:

$$P(\omega; 0) = \frac{e^2 \omega v(0)_z}{c^2} \left\{ \left[(\sin \theta_0) J_0(\xi_0) \right]^2 + \left[(v_\perp(0)/v_z(0)) J_1(\xi_0) \right]^2 \right\}, \quad (2a)$$

$$\cos \theta_0 = \frac{1}{n(\omega)\beta_z(0)}; \quad (2b)$$

$$\begin{aligned} P(\omega; a) = & \frac{e^2 v_z(a) \omega}{c^2} \left\{ \sin^2 \theta_a J_a^2(\xi_a) \right. \\ & + (v_\perp(a)/2v_z(a))^2 \left[\cos^2 \theta_a (J_{a+1}(\xi_a) + J_{a-1}(\xi_a))^2 + (J_{a+1}(\xi_a) - J_{a-1}(\xi_a))^2 \right] \\ & \left. - (v_\perp(a)/v_z(a)) \sin \theta_a \cos \theta_a J_a(\xi_a) [J_{a+1}(\xi_a) + J_{a-1}(\xi_a)] \right\}, \quad (3a) \end{aligned}$$

$$\cos \theta_a = \frac{1}{n(\omega)\beta_z(a)} - \frac{a\omega_c}{n(\omega)\omega\beta_z(a)}, \quad a = 1, 2, 3, \dots; \quad (3b)$$

$$\begin{aligned}
P(\omega; -b) = & \frac{e^2 v_z(-b) \omega}{c^2} \left\{ \sin^2 \theta_{-b} J_b^2(\xi_{-b}) \right. \\
& + \left(v_{\perp}(-b)/2v_z(-b) \right)^2 \left[\cos^2 \theta_{-b} \left(J_{b+1}(\xi_{-b}) + J_{b-1}(\xi_{-b}) \right)^2 + \left(J_{b+1}(\xi_{-b}) - J_{b-1}(\xi_{-b}) \right)^2 \right] \\
& + \left(v_{\perp}(-b)/v_z(-b) \right) \sin \theta_{-b} \cos \theta_{-b} J_b(\xi_{-b}) \left[J_{b+1}(\xi_{-b}) + J_{b-1}(\xi_{-b}) \right] \left. \right\}, \quad (4a)
\end{aligned}$$

$$\cos \theta_{-b} = \frac{1}{n(\omega) \beta_z(-b)} + \frac{b \omega_c}{n(\omega) \omega \beta_z(-b)}, \quad b = 1, 2, 3, \dots \quad (4b)$$

The notation is such that

$$\xi_l = \frac{n(\omega) \omega R_l \sin \theta_l}{c}, \quad R_l = \frac{v_{\perp}(l)}{\omega_c}, \quad \omega_c = \frac{eB}{M \gamma(l) c}, \quad l = 0, a, -b, \quad (5)$$

where, in general, R denotes the electron gyro-radius, M the mass of the electron, and $\gamma(l)$ is the electron relativistic factor. It is relations (2b), (3b), and (4b), of course, that kinematically define the HCE, harmonic radiation below, and above the helical Čerenkov threshold, respectively. For the sake of comparison, we also list the power spectrum of the ordinary Čerenkov effect (no magnetic field present; see, for example Šoln [1992]):

$$P(\omega; C) = \frac{e^2 \omega v_c}{c^2} \sin^2 \theta_c; \quad \cos \theta_c = \frac{1}{n(\omega) \beta_c}, \quad (6)$$

where v_c is the electron velocity and θ_c is the radiation angle.

2. NUMBER OF PHOTONS PER UNIT PATH LENGTH IN THE VISIBLE SPECTRUM

From now on, we shall assume that index of refraction n varies very slowly in the visible wavelength interval of interest; that is, for all practical purposes, n is independent of ω .

We start with the expression for the number spectrum per unit path length

$$\frac{d^2 N_T(\omega; l)}{dL d\omega} = \frac{P(\omega; l)}{\hbar \omega v_z(l)} \equiv N(\omega; l), \quad (7)$$

where $l = 0, a, -b$, and, by definition, $l = C$ represents the ordinary Čerenkov effect when v_z is replaced by v_c . Next, the number of photons per unit path length that are emitted into angular frequency interval $d\omega$ and the wavelength interval $d\lambda$ are given, respectively, as

$$\frac{d^2 N_T(\omega; l)}{dL d\omega} d\omega = N(\omega; l) d\omega = \tilde{N}(\lambda; l) d\lambda; \quad (8a)$$

$$\tilde{N}(\lambda; l) = \frac{P(\omega(\lambda); l)}{\hbar \lambda v_z(l)}. \quad (8b)$$

Because of $|d\omega/d\lambda| = \omega/\lambda$ (radiation is observed in a vacuum), we also have this very important relation,

$$N(\omega; l) \omega = \tilde{N}(\lambda; l) \lambda. \quad (9)$$

At this point, we introduce the "natural finite" wavelength and angular frequency intervals:

$$\Delta\lambda = \lambda_2 - \lambda_1 = \lambda_0, \lambda_{1,2} = \frac{\sqrt{5} \mp 1}{2} \lambda_0, \lambda_0 = \frac{\lambda_1 + \lambda_2}{\sqrt{5}}, \quad (10a)$$

$$\Delta\omega = \omega_1 - \omega_2 = \omega_0, \omega_{1,2} = \frac{\sqrt{5} \pm 1}{2} \omega_0, \omega_0 = \frac{\omega_2 + \omega_1}{\sqrt{5}}, \quad (10b)$$

where λ_0 and ω_0 are the center wavelength and center angular frequency, respectively. One can easily see that these definitions are numerically very close to what one uses in experiments (Martin and Shaw

1993). Furthermore, one also has that $\Delta\omega = 2\pi c/\Delta\lambda$. For latter references, we notice that to a good approximation $\sqrt{5} \equiv (e + 1)/(e - 1)$, which will be useful later. However, one can also introduce the "infinitesimal" wavelength and angular frequency intervals:

$$\delta\lambda = \lambda_0, \quad (11a)$$

$$\delta\omega = \omega_0. \quad (11b)$$

It will be seen that for the ordinary and helical Čerenkov effects the calculated (dN_T/dL) 's for radiation falling within either $\Delta\lambda$ or $\delta\lambda$ are practically the same.

For the ordinary and HCE, one has from relations (8) that the total number of photons per unit path length emitted within wavelength and angular frequency intervals $\Delta\lambda$ or $\Delta\omega$ to be, respectively

$$\frac{dN_T(\lambda_0, \Delta\lambda; l = 0, C)}{dL} = \int_{\lambda_1}^{\lambda_2} \tilde{N}(\lambda; l = 0, C) d\lambda, \quad (12a)$$

$$\frac{dN_T(\omega_0, \Delta\omega; l = 0, C)}{dL} = \int_{\omega_2}^{\omega_1} N(\omega; l = 0, C) d\omega. \quad (12b)$$

Of course, (12a) = (12b). Next we estimate the contributions from harmonics to dN_T/dL within $\Delta\omega$ or $\Delta\lambda$ intervals. Since harmonics occur at sharp frequencies (wavelengths), it is sufficient to estimate them for radiation that falls within infinitesimal $\delta\omega$ ($\delta\lambda$). Of course, $\delta\omega$ ($\delta\lambda$) is supposed to be within $\Delta\omega$ ($\Delta\lambda$). Having said that, consistent with relations (11), we have that for any harmonic index l

$$\frac{dN_T(\lambda_0, \delta\lambda; l)}{dL} = \tilde{N}(\lambda_0; l) \lambda_0 = \frac{dN_T(\omega_0, \delta\omega; l)}{dL} = N(\omega_0, l) \omega_0. \quad (13a,b)$$

With these preliminaries, we first deal with the ordinary Čerenkov effect. Consistent with relations (6), (8), (10), and (12), we obtain

$$\frac{dN_T(\lambda_0, \Delta\lambda; C)}{dL} = 2\pi\alpha\sin^2\theta_c \left(\frac{1}{\lambda_1} - \frac{1}{\lambda_2} \right) = \tilde{N}(\lambda_0, C)\lambda_0 = \frac{dN_T(\lambda_0, \delta\lambda; C)}{dL}, \quad (14a)$$

$$\frac{dN_T(\omega_0, \Delta\omega; C)}{dL} = \frac{\alpha}{c} \sin^2\theta_c (\omega_1 - \omega_2) = N(\omega_0, C)\omega_0 = \frac{dN_T(\omega_0, \delta\omega; C)}{dL}. \quad (14b)$$

As we see, expressions (14a) and (14b) are exactly in the forms of relations (13a) and (13b), respectively.

To address the HCE and harmonic radiation, we first rewrite the argument of the Bessel Functions from (5) as

$$\xi_l = (\pi\gamma(l)\beta_\perp(l)\sin\theta_l) \frac{\omega(Mc/e)}{B}. \quad (15)$$

In the visible spectrum, we have that typically $\omega \cong 4 \times 10^{15} \text{s}^{-1}$. With $(Mc/e) \cong 5.56 \times 10^{-12} \text{sT}$, we have that $\xi_l \cong (\pi\gamma(l)\beta_\perp(l)\sin\theta_l) 2 \times 10^4 \text{T/B}$ is going to be very large for "moderate" magnetic fields, $B \leq 100 \text{ T}$. Hence, we can utilize the asymptotic expressions for Bessel functions (Arfken 1985):

$$8x \gg 4l^2 - 1: J_l^2(x) \cong \frac{1}{\pi x} [1 + (-1)^l \sin 2x]. \quad (16)$$

Since in the HCE only $l = 0, 1$ come, we are allowed to use the asymptotic expressions for Bessel functions in form (16). With relations (2a), (7), (8), and (9), we obtain

$$N(\omega; l=0) = N_{eff}(\omega; l=0) + N_{osc}(\omega; l=0), \quad (17a)$$

$$\tilde{N}(\lambda; l=0) = \tilde{N}_{eff}(\lambda; l=0) + \tilde{N}_{osc}(\lambda; l=0); \quad (17b)$$

$$N_{eff}(\omega; l=0) = \frac{\alpha[(n\beta(0))^2 - 1]}{c\pi\xi_0(n\beta_z(0))^2}, \quad (18a)$$

$$N_{osc}(\omega; l=0) = \frac{\alpha \sin 2\xi_0 [(n\beta_z(0))^2 - (n\beta_z(0))^2 - 1]}{c\pi\xi_0(n\beta_z(0))^2}, \quad (18b)$$

$$\tilde{N}_{eff,osc}(\lambda; l=0) = \frac{2\pi c}{\lambda^2} N_{eff,osc}(\omega(\lambda); l=0). \quad (18c)$$

It is evident that generally $|N_{osc}| < |N_{eff}|$. Furthermore, as λ varies from λ_1 through λ_0 to λ_2 , N_{osc} experiences rapid oscillations in the variable ξ_0 ; as such, its contribution in the integral (12b) is negligible. Therefore, we obtain

$$\begin{aligned} \frac{dN_T(\lambda_0, \Delta\lambda; l=0)}{dL} &= \tilde{N}_{eff}(\lambda_0; l=0) \lambda_0 \ln(\lambda_2/\lambda_1) \\ &\equiv \tilde{N}_{eff}(\lambda_0; l=0) \lambda_0 = \frac{dN_T^{eff}(\lambda_0, \delta\lambda; l=0)}{dL}, \end{aligned} \quad (19a)$$

$$\begin{aligned} \frac{dN_T(\omega_0, \Delta\omega; l=0)}{dL} &= N_{eff}(\omega_0; l=0) \omega_0 \ln(\omega_1/\omega_2) \\ &\equiv N_{eff}(\omega_0; l=0) \omega_0 = \frac{dN_T^{eff}(\omega_0, \delta\omega; l=0)}{dL}, \end{aligned} \quad (19b)$$

where, of course, (19a) = (19b) term by term. Next, in relations (19), because of $|N_{osc}| < |N_{eff}|$, one may write $N_{eff}(l=0) + N_{osc}(l=0)$ instead of just $N_{eff}(l=0)$ and still obtain a good estimate for $dN_T(l=0)/dL$; this we then denote as $dN_T^{est}(l=0)/dL$.

Examples (19) suggest, then, that for harmonic radiation above and below the HCE threshold, rather than going through tedious calculations, one may simply estimate the number of photons per unit length emitted within $\Delta\omega$ or $\Delta\lambda$, respectively, as

$$\frac{dN_T^{est}(\omega_0, \Delta\omega; l=a_0, -b_0)}{dL} \equiv \frac{dN_T(\omega_0, \delta\omega; l=a_0, -b_0)}{dL} = N(\omega_0; l=a_0, -b_0)\omega_0, \quad (20a)$$

$$\frac{dN_T^{est}(\lambda_0, \Delta\lambda; l=a_0, -b_0)}{dL} \equiv \frac{dN_T(\lambda_0, \delta\lambda; l=a_0, -b_0)}{dL} = \tilde{N}(\lambda_0; l=a_0, -b_0)\lambda_0, \quad (20b)$$

where again (20a) = (20b) term by term and a_0 and b_0 are harmonic indices associated with the central angular radiation frequency ω_0 according to relations (3) and (4), respectively.

3. APPLICATIONS

On general grounds, it follows from relations (2) to (5) that as B becomes larger and/or $\beta_\perp(l)$ becomes smaller, the HCE will dominate over the harmonic radiation. In fact, for $\beta_\perp \neq 0$, and $B \rightarrow 0$, the HCE will gradually disappear, while the radiation into harmonics will become more dominant. At $B = 0$, all the harmonic radiation will sum up into the ordinary Čerenkov radiation, which, however, can also be calculated directly (Šoln 1992).

Let us go back to the nonmonoenergetic beam. Here, by definition, the estimated overall number of photons per unit path length is the superposition of the emitted total number of HCE photons per unit path length and the estimated total number of photons per unit path length into the harmonics above and below the HCE threshold. We shall assume that as one goes from one electron to another in the beam that $\beta_\perp(a) = \beta_\perp(-b) = \beta_\perp(0) = 0.3$, where $l = 0, a, -b$ are indices associated with the HCE and harmonic radiation below and above the HCE threshold, respectively. Their constancy is, of course, an average value for β_\perp across the beam; it is justified by the fact that β_\perp does not enter into the definition of radiation frequency. Hence, the nonmonoenergetic beam quality will be specified by $\beta_z(l)$ which approximately varies between 0.92 and 0.94 with $\beta_z(0) = 0.934$ corresponding to the HCE (the energies of individual electrons are in the 2–3-MeV range). The helical motion of electrons is maintained through silica aerogel as a medium (the index of refraction in the visible spectrum is $n = 1.075$) by the magnetic

field of $B = 10$ T. The radiation angular frequency in all three cases is simply the central angular radiation frequency in the visible portion of the spectrum, $\omega_0 = 3.77 \times 10^{15} \text{s}^{-1}$. Specifically, using relations (19), (20), (7), and (2) to (5), we have the following expressions for the number of photons per unit path length for each case:

$$l = 0, \beta_z(0) = 0.9340, \cos\theta_0 = 1/(n\beta_z(0)) = 0.9960, \gamma(0) = 5.1541:$$

$$\frac{dN_T(\omega_0, \Delta\omega; l=0)}{dL} \equiv 0.104 \text{ cm}^{-1}; \quad (21a)$$

$$l = a_0 \equiv 120, \beta_z(a_0) = 0.9206, \cos\theta_{a_0} = \cos\theta_0, \gamma(a_0) = 4:$$

$$\frac{dN_T^{est}(\omega_0, \Delta\omega; l=a_0)}{dL} \equiv 0.205 \text{ cm}^{-1}; \quad (21b)$$

$$l = -b_0 \equiv -71, \beta_z(-b_0) = 0.9393, \cos\theta_{-b_0} = \cos\theta_0, \gamma(-b_0) = 6:$$

$$\frac{dN_T^{est}(\omega_0, \Delta\omega; l=-b_0)}{dL} \equiv 0.120 \text{ cm}^{-1}. \quad (21c)$$

At the end of the $L = 10$ -cm path length in the silica aerogel, the number of photons generated by an electron in the radiation angular frequency interval $\Delta\omega$ with the central radiation angular frequency $\omega_0 = 3.77 \times 10^{15} \text{s}^{-1}$ and at the central angle θ_0 are, respectively

$$N_T(\omega_0, \Delta\omega; l=0) = 1.04, \quad (22a)$$

$$N_T^{est}(\omega_0, \Delta\omega; l=a_0) = 2.05, \quad (22b)$$

$$N_T^{est}(\omega_0, \Delta\omega; l=-b_0) = 1.2, \quad (22c)$$

where the parameters correspond to relations (21a,b,c).

We can take a simple average over $l = 0, a_0, -b_0$ of relations (21a,b,c) yielding

$$\frac{dN_T^{est}(\omega_0, \Delta\omega; average)}{dL} \equiv 0.143 \text{ cm}^{-1}. \quad (23)$$

Comparing this expression with (21a), we see that the harmonic emission actually enhanced the helical Čerenkov radiation. Finally, from relation (23), we have

$$L = 10 \text{ cm}, B = 10 \text{ T} : N_T^{est}(\omega_0, \Delta\omega; average) = 1.43, \quad (24)$$

so that an electron beam with, say, 10^8 electrons in cm^3 will generate about 1.4×10^8 photons in cm^3 at the end of a 10-cm path length in the silica aerogel. This can be observed experimentally.

Recently, a great deal of progress has been made in achieving magnets whose fields would reach 100 T (Boebinger, Passner, and Bevk 1995). With $B = 100 \text{ T}$ in relations (21a,b,c), the harmonic numbers and the number of photons per unit path length become, respectively: 0, 1.039 cm^{-1} ; 12, 2.166 cm^{-1} ; and $-7, 0.408 \text{ cm}^{-1}$. One notices that only for the HCE, $l = 0$, the number of photons per unit path length scales linearly with B .

These examples show explicitly that the HCE with a nonmonoenergetic electron beam could be a vehicle for detecting a magnetic field in the hostile electromagnetic medium.

INTENTIONALLY LEFT BLANK.

4. REFERENCES

Arfken, G. Mathematical Methods for Physicists. Third Edition. New York: Academic Press, p. 620, 1985.

Boebinger, G., A. Passner, and J. Bevk. "Building World-Record Magnets." Scientific American, pp. 58–66, June 1995.

Martin, B. R., and G. Shaw. Particle Physics. New York: John Wiley & Sons, pp. 69–71, 1993.

Šoln, J. Phys. Rev., vol. A46, p. 5138–5148, 1992.

INTENTIONALLY LEFT BLANK.

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	DEFENSE TECHNICAL INFO CTR ATTN DTIC DDA 8725 JOHN J KINGMAN RD STE 0944 FT BELVOIR VA 22060-6218

1	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TA 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

3	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TL 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

1	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TP 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

ABERDEEN PROVING GROUND

5	DIR USARL ATTN AMSRL OP AP L (305)
---	---------------------------------------

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>	<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
1	OSD OUSD AT STRT TAC SYS ATTN DR SCHNEITER 3090 DEFNS PENTAGON RM 3E130 WASHINGTON DC 20301-3090	1	ARMY RESEARCH LABORATORY ATTN AMSRL ST DR FRASIER 2800 POWDER MILL RD ADELPHI MD 20783-1197
1	ODDRE AT ACQUISITION AND TECH ATTN DR GONTAREK 3080 DEFENSE PENTAGON WASHINGTON DC 20310-3080	1	ARMY RESEARCH LABORATORY ATTN AMSRL SL DR WADE WSMR NM 88002-5513
1	ASST SECY ARMY RESEARCH DEVELOPMENT ACQUISITION ATTN SARD ZD RM 2E673 103 ARMY PENTAGON WASHINGTON DC 20310-0103	1	ARMY RESEARCH LABORATORY ATTN AMSRL SL E MR MARES WSMR NM 88002-5513
1	ASST SECY ARMY RESEARCH DEVELOPMENT ACQUISITION ATTN SARD ZP RM 2E661 103 ARMY PENTAGON WASHINGTON DC 20310-0103	1	ARMY TRADOC ANL CTR ATTN ATRC W MR KEINTZ WSMR NM 88002-5502
1	ASST SECY ARMY RESEARCH DEVELOPMENT ACQUISITION ATTN SARD ZS RM 3E448 103 ARMY PENTAGON WASHINGTON DC 20310-0103	1	ARMY TRNG & DOCTRINE CMND ATTN ATCD B FT MONROE VA 23651
1	ASST SECY ARMY RESEARCH DEVELOPMENT ACQUISITION ATTN SARD ZT RM 3E374 103 ARMY PENTAGON WASHINGTON DC 20310-0103	1	<u>ABERDEEN PROVING GROUND</u>
1	UNDER SEC OF THE ARMY ATTN DUSA OR RM 2E660 102 ARMY PENTAGON WASHINGTON DC 20310-0102	1	CDR USATECOM ATTN: AMSTE-TA
1	ASST DEP CHIEF OF STAFF OPERATIONS AND PLANS ATTN DAMO FDZ RM 3A522 460 ARMY PENTAGON WASHINGTON DC 20310-0460	2	CDR USAMSAA ATTN: AMXSY-ST AMXSY-D
1	DEPUTY CHIEF OF STAFF OPERATIONS AND PLANS ATTN DAMO SW RM 3C630 400 ARMY PENTAGON WASHINGTON DC 20310-0400	4	DIR USARL ATTN: AMSRL-SL, J WADE (433) AMSRL-SL-I, M STARKS (433) AMSRL-SL-C, W HUGHES (E3331) AMSRL-SL-B, P DEITZ (328)
		1	CDR CBDCOM ATTN: TECHNICAL LIBRARY BLDG E3330
		1	DIR CBIAC BLDG E3330, RM 150

NO. OF COPIES	ORGANIZATION
2	DIRECTOR US ARMY RESEARCH LABORATORY ATTN AMSRL EP M V GELNOVATCH AMSRL EP EF M DUTTA FT MONMOUTH NJ 07703-5601
1	DIRECTOR US ARMY RESEARCH LABORATORY ATTN AMSRL SS V DEMONTE ADELPHI MD 20783-1197
1	DIRECTOR US ARMY RESEARCH LABORATORY ATTN AMSRL OP TL WATERTOWN MA 02172-0001
2	DIRECTOR US ARMY RESEARCH LABORATORY ATTN AMSRL SS IC P EMMERMAN R WINKLER ADELPHI MD 20783-1197
1	DIRECTOR US ARMY BALLISTIC MIS DEFNS SYS CMND ATTN ADVANCED TECHNOLOGY CENTER PO BOX 1500 HUNTSVILLE AL 35807-3801
2	COMMANDER US ARMY STRATEGIC DEFENSE COMMAND ATTN CSSD SL C CSSD SL S HUNTSVILLE AL 35807-3801
1	COMMANDER CECOM R&D TECHNICAL LIBRARY ATTN ASQNC ELC IS L R FORT MONMOUTH NJ 07703-5000
1	COMMANDER US ARMY RESEARCH OFFICE ATTN TECHNICAL LIBRARY PO BOX 12211 RESEARCH TRIANGLE PARK NC 27709-2211
1	COMMANDER US ARMY LOGISTIC MGMT CENTER ATTN DEFENSE LOGISTICS STUDIES FORT LEE VA 23801
1	COMMANDANT US ARMY CMND AND GEN STAFF COLLEGE FORT LEAVENWORTH KS 66027

NO. OF COPIES	ORGANIZATION
1	COMMANDER US ARMY NGIC ATTN AMXST MC 3 220 SEVENTH STREET NE CHARLOTTESVILLE VA 22901-5396
8	DIR, USARL ATTN AMSRL SL E G MARES AMSRL SL EA R SHELburne AMSRL SL EG J PALOMO AMSRL SL EM R FLORES AMSRL SL EP D ALVAREZ AMSRL SL ES T ATHERTON AMSRL SL EV DR K MORRISON AMSRL SL CA R SUTHERLAND WHITE SANDS MISSILE RANGE NM 88002-5513 <u>ABERDEEN PROVING GROUND, MD</u>
18	DIR, USARL ATTN: AMSRL-SL, J. SMITH AMSRL-SL-BA, DR. J. WALBERT J. HANES R. KUNKEL L. ROACH AMSRL-SL-BS, D. BELY DR. T. KLOPCIC AMSRL-SL-BV, L. MORRISSEY W. MERMAGEN, JR. AMSRL-SL-I, R. REITZ D. BASSETT A. YOUNG M. VOGEL E. PANUSKA DR. D. HASKELL DR. J. FEENEY R. ZIGLER AMSRL-WT, DR. H. E. BRANDT
1	CDR, USAATC ATTN: STECS
1	DIR, ERDEC ATTN: SCBRD-RT

NO. OF COPIES	ORGANIZATION
------------------	--------------

58	DIR, USARL ATTN: AMSRL-SL-C, J. SEIGH MAJ. JEROME GILMAN AMSRL-SL-CM, D. FARENWALD B. RUTH L. DAVIS L. D'ELICIO R. JOLLIFFE D. SLOOP R. TRYTUS R. ZUM BRUNNEN E. FIORAVANTE M. MAR DR. J. SOLN (30 CYS) AMSRL-SL-CO, D. BAYLOR R. PROCHAZKA AMSRL-SL-CS, J. BEILFUSS T. FLORY B. SMITH DR. M. SMITH J. FRANZ T. MAK J. NEALON R. PARSONS M. KAUFMAN D. MANYAK R. POLIMADEI M. BUMBAUGH J. CAPOBIANCO D. DAVIS
----	--

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. ARL Report Number ARL-TR-947 Date of Report January 1996

2. Date Report Received _____

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

4. Specifically, how is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. _____

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) _____

CURRENT
ADDRESS

Organization

Name

Street or P.O. Box No.

City, State, Zip Code

7. If indicating a Change of Address or Address Correction, please provide the Current or Correct address above and the Old or Incorrect address below.

OLD
ADDRESS

Organization

Name

Street or P.O. Box No.

City, State, Zip Code

(Remove this sheet, fold as indicated, tape closed, and mail.)
(DO NOT STAPLE)

DEPARTMENT OF THE ARMY


OFFICIAL BUSINESS

BUSINESS REPLY MAIL

FIRST CLASS PERMIT NO 0001,APG,MD

POSTAGE WILL BE PAID BY ADDRESSEE

DIRECTOR
U.S. ARMY RESEARCH LABORATORY
ATTN: AMSRL-SL-CM
ABERDEEN PROVING GROUND, MD 21010-5423



NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

